

There are no translations available.

**C.81-Voltage-gated sodium Na<sup>+</sup> channel allostery as the basis for the scorpion venom model of AIDS:Molecular homology between between spider toxin and HIV-2, and scorpion toxin and HIV-1 envelope gp41 sequence SWSNKS**

Abstract accepté pour EMBO Conference on Allosteric interactions in cell signaling and regulation(Institut Pasteur Paris,May 14-17, 2013 )

**G.M.K. Tran**

31 avenue du bois 92290 Châtenay-Malabry, France ; Association « Positifs », BP 230, 75865Paris cedex 18, France

**Objective** : Allosteric differences explain why spider Atrax toxin differs slightly from scorpion Androctonus AaH2, despite the fact they bind to the same Na<sup>+</sup> channel receptor site 3 (Little MJ, 1998). We found a molecular mimicry between HIV-1 envelope gp 120 V3 and V2 loops, but nothing between scorpion and HIV-2, which is less virulent than HIV-1.

**Methods** : Amino acid and three-dimensional (3D) structure (Pallaghy PK, 1997) comparison between

HIV-2/SIV and the arthropod family, conotoxin and sea anemone toxin, all ligands of the Na<sup>+</sup> channel.

**Results** : 1°) There is a match between lethal Australian spider Atrax Robustus toxin and HIV-2, but only in 3D, although the structures differed: Spider key residues (4-KR-5/M18/K19/Y22/23-AWY-25/Q27/C31) superimpose on HIV-2 residues (180-KK-181/M172/K173/Y183/186-AWY-188/Q190/C194).

Written by GMK Tran

Friday, 12 April 2013 14:26 - Last Updated Friday, 14 June 2013 12:59

---

2°) We confirm the scorpion model of HIV-1, by aligning scorpion toxin COOH-terminus with

HIV-1/chimpanzee CPZ.CD90.ANT gp41 sequence SWSNKS, or 3S, a potential therapeutic AIDS vaccine (Vieillard V, 2008).WSNKS is deleted in HIV-2/SIV-AGM (explaining why African Green Monkey AGM don't get sick):

scorpion chimera C...KLAC.CY-SVP-WNPTWS-RSNTCGKK-carboxy-terminus

HIV-1/SIVchimp C..-KLV..CYTSVPDWVPSWSNKSQTCAKN

**Conclusions** : Na<sup>+</sup> channel has many sites, the site 3 binding scorpion and spider toxins, namely respectively HIV-1 and HIV-2 envelope; their different degree of virulence is explained by the different allosteric modifications between scorpion and spider toxins. Another implication is to ameliorate the clinical results obtained with a Na<sup>+</sup> channel ligand, omega 3, which is effective and non toxic in AIDS treatment. By combining many ligands